

Taking Your Hepatitis C Therapy:

Zepatier™ with or without Ribavirin

Elbasvir/Grazoprevir = Zepatier™



- ▶ Dose: 1 tablet (50/100 mg) daily, taken at the same time each day
- ▶ You can take with or without food
- ▶ Store at room temperature between 59°F and 86°F
- ▶ Keep tablets in the original package until use to protect from moisture

Do You Need to Take Ribavirin? Yes No








- ▶ Dose: 200 mg capsules or tablets; your dose may change during the course of treatment
- ▶ Take with food to minimize stomach upset such as nausea and vomiting
- ▶ Do not consider pregnancy until at least 6 months after treatment for either partner has ended
- ▶ Store at room temperature (< 86°F) away from direct sunlight

Daily Medication Schedule

Start Date: _____ End Date: _____

Expected Treatment Duration: 12 weeks 16 weeks

Morning Time: ____ am	Evening Time: ____ pm
 Take 1 Elbasvir/Grazoprevir tablet daily	
 Take ____ Ribavirin with food 	 Take ____ Ribavirin with food 

Note: Elbasvir/Grazoprevir and ribavirin can be taken together or at different times, but keep to the same time of day for each drug.

If You Miss a Dose:

Elbasvir/Grazoprevir: Zepatier™

- ▶ If you miss a dose, TAKE THE MISSED DOSE THE SAME DAY as soon as you remember; take your next dose of Elbasvir/Grazoprevir at your regular time the next day
- ▶ Do not take more than 1 tablet of Elbasvir/Grazoprevir in a day
- ▶ DO NOT take two doses at the same time to make up for a missed dose
- ▶ Note: If you miss multiple doses, call your prescriber/clinic listed below

Ribavirin

- ▶ If you miss more than 1-2 days of Ribavirin, call your prescriber/clinic listed below

Prescriber/Clinic Contact: _____

For Refills: _____



U.S. Department of Veterans Affairs

Veterans Health Administration
Patient Care Services